

<b>Order Anseriformes</b> <b>Whistling-Ducks, Geese, Swans, and Ducks</b>	Scaup, Greater	10A-4B Mu WRu
Whistling-Duck, Black-bellied*	Lesser	10D-4D+ Mc WRc
Fulvous*	Scoter, Surf	10B-5B Mr WRvr
Goose, Greater White-fronted	White-winged	10D-5D+ Mr Wvr
Snow	Black	10D-3C Mvr WRvr
Ross's	Duck, Long-tailed	11A-4A+ Mr Wvr
Brant	Bufflehead	11A-4C+ WRc
Goose, Barnacle	Goldeneye, Common	11B-4C WRLfc
Cackling	Barrow's	11A-3D+ WRu (Lfc)
Canada*	Merganser, Hooded*	P c SRLu WRfc
Swan, Mute	Common	10A-3C+ Wvr (n)
Trumpeter	Red-breasted	10D-4D+ Mu Wvr
Tundra	Duck, Ruddy°	10A-4C+ Mc WRc SRr
Duck, Wood*	<b>Order Galliformes</b> <b>Turkey and Quail</b>	
Gadwall	Bobwhite, Northern*	P Lu
Wigeon, Eurasian	Grouse, Ruffed	extirpated
American	Prairie-Chicken, Greater	extirpated
Duck, American Black	Turkey, Wild*	P u
Mallard*	<b>Order Gaviformes</b> <b>Loons</b>	
Duck, Mottled*	Loon, Red-throated	10D-4A Mvr WRr
Teal, Blue-winged*	Pacific	11A-4A+ Mr WRr
Cinnamon	Common	10A-5D+ Mfc WRLc
Shoveler, Northern	Yellow-billed	FWVacc (2, 2010)
Pintail, Northern	<b>Order Podicipediformes</b> <b>Grebes</b>	
Garganey	Grebe, Least	FVacc (1, 1984)
Teal, Green-winged	Pied-billed*	P WRc SRLu
Canvasback	Horned	10A-4A+ WRfc
Redhead	Red-necked	11D-4B WVvr
Duck, Ring-necked	Eared	9A-5A+ Mu WRr (Lu se)
Tufted	Western	10B-3D+ Mr Wvr

<b>Order Ciconiiformes</b> <b>Storks</b>		Osprey*	P Mfc SRLr WRr (s)
Stork, Wood	6B-10B+ FVu (s, e)	Kite, Swallow-tailed°	4A-9D Mvr SRLvr
<b>Order Suliformes</b> <b>Frigatebirds, Boobies, Cormorants, and Anhinga</b>		White-tailed	3A-4D+ SpWVvr
Frigatebird, Magnificent	9B-10D+ Sa FVvr	Mississippi*	4C-9C+ SRc (Lu wc, nw)
Booby, Brown	Vocc (4, 2013)	Eagle, Bald*	P WRc SRu
Gannet, Northern	SVacc (1, 2005)	Harrier, Northern°	8B-5B+ Mfc WRfc
Cormorant, Neotropic°	5C-12D+ Mr SRLr (sw, se) WVvr	Hawk, Sharp-shinned*	P c Mfc WRu SRr (n)
Double-crested*	P Mc SRLu WRc	Cooper's*	P WRfc SRLu
Anhinga*	3C-11C+ SRLu WVvr (s)	Goshawk, Northern	11A-1B WVvr
<b>Order Pelicaniformes</b> <b>Pelicans, Herons, Ibises, and Spoonbills</b>		Hawk, Red-shouldered*	P fc
Pelican, American White	P Mfc SRLu WRfc	Broad-winged*	3D-10B+ Mfc SRfc
Brown	3C-11B+ Vr (ce, s)	Swainson's*	3C-11C+ Mu (w) SRLr (nw)
Bittern, American°	3A-5D 9B-12D+ Mu WVvr	Red-tailed*	P c
Least*	4A-10A+ Mr SRr (Lu s, e)	Rough-legged	10D-3C WVvr
Heron, Great Blue*	P c	Ferruginous	10D-2D WVvr
Egret, Great*	P SRLc WRfc (ce, s)	Eagle, Golden	9B-4B WRr
Snowy*	3B-11C+ Mfc SRLfc WVvr	<b>Order Gruiformes</b> <b>Rails, Gallinules, and Cranes</b>	
Heron, Little Blue*	3B-11C+ Mc SRLc WVvr	Rail, Yellow	9A-5A Mvr WVocc
Tricolored*	4A-10C+ Mr SRLu (s) FVu (s) WVvr (sw)	Black	9B-10B FMvr
Egret, Cattle*	3B-11B+ Mfc SRc WVvr (s)	King*	P Mr SRLr (ce, s)
Heron, Green*	3B-10D+ Mc SRc WVvr (s)	Virginia	4A-5D 8D-2D+ MLu WRLr
Night-Heron, Black-crowned*	3B-12B+ Mu SRLu WVvr	Sora	4A-5C 9A-12D+ Mfc WRr (ce, s)
Yellow-crowned*	3D-11A+ Mu SRLu	Gallinule, Purple*	4B-10C SRLr (ce, s)
Ibis, White*	3D-11A+ SRLc (s, sw) FVLc (s, ce) WRvr (sw)	Common*	4B-12D+ Mr (nw) SRLu (s) WRLr (sw)
Glossy°	4B-5D 7C-9D+ Mr	Coot, American*	P Mc WRc SRLr
White-faced	4B-6A 8A-11D+ Mr WVvr (se)	Crane, Sandhill	10A-5A+ Mr WVr WRLu (ne, e)
Spoonbill, Roseate	7A-10D+ FVu (ce, s)	Whooping	extirpated
<b>Order Accipitriformes</b> <b>Vultures, Kites, Eagles, and Hawks</b>		<b>Order Charadriiformes</b> <b>Plover, Sandpipers, Gulls, and Terns</b>	
Vulture, Black*	P fc (w) r (e)	Stilt, Black-necked*	3D-10D+ SRfc (ec, se)
Turkey*	P c	Avocet, American	4A-11D+ SpMu FMfc

Plover, Black-bellied	3C-5D 8B-11D+ SpMu FMu	Sandpiper, Buff-breasted	4A-5D 7D-10A+ SpMvr FMu
Golden-Plover, American	3A-5D 8C-11B+ SpMc FMu	Pectoral	3A-6B 6D-11D+ Mc
Plover, Snowy	FSpVocc (5, 2014)	Semipalmated	4A-6B 7B-11A+ Mc
Wilson's	SpFVocc (3, 1998)	Western	4B-6A 7A-12D+ SpMu FMfc WRvr
Semipalmated	4B-5C 7B-9D+ Mc	Dowitcher, Short-billed	4D-6A 7D-10A Mu
Piping	4B-5C 7B-10B SpMvr FMr	Long-billed	3A-5B 7A-12D+ Mc WRLr
Killdeer*	P c	Snipe, Wilson's	8A-5D+ Mc WRfc
Plover, Mountain	WVacc (1, 1951)	Woodcock, American*	P SpMfc SRr FMu WRu
Sandpiper, Spotted°	3D-6B 7A-12D+ Mc Wvr	Phalarope, Wilson's	3C-5D 7A-10D Mfc
Solitary	3D-5B 6B-12D+ Mfc	Red-necked	5B-6A 7B-11B SpMvr FMr
Yellowlegs, Greater	3B-5D 7A-12D+ Mc WRu	Red	8D-11D+ FMvr
Willet	3D-5D 6C-9C+ Mu	Jaeger, Pomarine	FVacc (1, 1983)
Yellowlegs, Lesser	2D-5D 6D-12D+ Mc WRu	Parasitic	9A-12B FVvr
Sandpiper, Upland	3D-5C 6D-9D+ Mu	Long-tailed	FVocc (4, 2000)
Curlew, Eskimo	extirpated	Murrelet, Long-billed	FVacc (1, 1980)
Whimbrel	5B-5D+ 9A-9B SpMvr FMacc	Kittiwake, Black-legged	11B-3A WVvr
Curlew, Long-billed	4B-4D 10B-12B Macc	Gull, Sabine's	9A-10C+ FMLr (sw, wc)
Godwit, Hudsonian	4B-5D 8C-10B SpMLu FMvr	Bonaparte's	10A-4D+ Mfc WRfc
Marbled	4B-5D 7C-10A SpMvr FMr	Black-headed	11C-1B WVvr
Turnstone, Ruddy	5B-6B 7C-10B Mr	Little	11D-3C+ WVvr
Knot, Red	5C-5D 8B-10A SpMacc FMvr	Laughing	7A-5B+ Vr
Ruff	3C-5B 8A-9B SpMvr FMacc	Franklin's	3B-6A 9C-12C+ SpMu FMfc
Sandpiper, Stilt	3D-6A 7A-11D+ Mfc	Ring-billed	8C-5A+ WRc
Curlew	FVacc (2, 2004)	California	12A-3B+ WVvr (wc)
Sanderling	4C-5D 7C-10D+ SpMr FMfc	Herring	8B-5C+ WRLu
Dunlin	3B-5D 10A-12D+ Mfc WRr	Thayer's	11D-3A WVvr (wc)
Sandpiper, Purple	FVacc (1, 1976)	Lesser Black-backed	10B-3B WVvr WRLr (wc)
Baird's	3B-6A 7D-11C SpMu FMfc	Glaucous	11D-3B WVvr
Least	7B-6B+ Mc WRu	Great Black-backed	WVacc (1, 2015)
White-rumped	4B-6C 8B-10D SpMfc FMvr	Tern, Sooty	Sa FVvr (9, 2014)

Tern, Bridled	Sa FVacc (2, 2008)	Owl, Short-eared	10C-4C WRLu
Least*	5A-9D+ Mu SRLu (Lc e)	Northern Saw-whet	11A-2D WVvr
Gull-billed	SpVacc (1, 2009)	<b>Order Caprimulgiformes</b> <b>Goatsuckers</b>	
Caspian	4A-6C 7A-10C+ SpMu FMfc	Nighthawk, Common*	4B-11A+ SRc
Black	4B-6B 6D-10C+ Mc	Chuck-will's-widow*	3D-9C SRc
Common	4D-5D 7A-10C+ SpMr FMu	Whip-poor-will, Eastern*	3D-9D SRfc (n, ce, wc)
Forster's	7B-5C+ Mfc WRLu (s)	<b>Order Apodiformes</b> <b>Swifts and Hummingbirds</b>	
Royal	Sa FVvr (6, 2012)	Swift, Chimney*	3B-10D SRc
Skimmer, Black	Sa FVvr (8, 2008)	White-throated	SpFWVacc (4, 2012)
<b>Order Columbiformes</b> <b>Pigeons and Doves</b>		Violetear, Green	6A-10A FVvr
Pigeon, Rock*	P c	Hummingbird, Magnificent	SVacc (1, 1993)
Collared-Dove, Eurasian*	P c	Ruby-throated*	3B-11B+ SRc
Pigeon, Passenger	extinct	Black-chinned	10D-3D WVvr
Dove, Inca*	P u Lfc (s, wc)	Anna's	10C-3B WVvr
Ground-Dove, Common	8C-5A Vr	Broad-tailed	FWVacc (4, 2006)
Dove, White-winged*	P Vr Lu (w)	Rufous	8B-4A FMr WVvr
Mourning*	P c	Calliope	FVacc (6, 2010)
<b>Order Cuculiformes</b> <b>Cuckoos, Roadrunners, and Anis</b>		Broad-billed	FWVacc (2, 2009)
Cuckoo, Yellow-billed*	4A-10C+ SRc	Buff-bellied	WVacc (1, 1993)
Black-billed°	4C-6D 8C-10C+ Mr SRvr (n)	<b>Order Coraciiformes</b> <b>Kingfishers</b>	
Roadrunner, Greater*	Lu (w, ce) Lr (e)	Kingfisher, Belted*	P c
Ani, Groove-billed	5C-5D 9C-11D+ SpVacc FVvr	<b>Order Piciformes</b> <b>Woodpeckers</b>	
<b>Order Strigiformes</b> <b>Owls</b>		Woodpecker, Lewis's	SVacc (1, 1967)
Owl, Barn*	P Lu (e) Lr (w)	Red-headed*	P fc
Screech-Owl, Eastern*	P fc	Red-bellied*	P c
Owl, Great Horned*	P fc	Sapsucker, Yellow-bellied	9B-5B+ WRc
Snowy	11B-1B WVvr	Woodpecker, Downy*	P c
Burrowing	9D-3D WVvr (e, ce)	Hairy*	P fc
Barred*	P fc	Red-cockaded*	P Lr (ec, se, wc)
Long-eared	11A-5D+ WRvr	Flicker, Northern*	P SRu WRfc

**Reporting Bird Sightings**

Ornithology has benefited a great deal from the contributions of amateurs over the years. Birders who faithfully report significant observations and make them a part of the permanent record help ornithologists better understand the intricacies of bird distribution and abundance. Although it isn't always obvious whether a sighting should be reported, a good rule to follow is "If in doubt, make the report." More specific instances in which a report should be made include:

- If a species appears outside of its normal season or geographical region, it should be reported.
- If unexpectedly large or small numbers of a certain species are encountered, that count of individuals should be reported.
- If an unusual or unlisted species is seen, report it by all means, but keep in mind that the greater the rarity the more thorough supporting documentation should be.

**Written and/or photographic documentation is required when reporting rarities or species observed out of season.** The Arkansas Bird Records Committee reviews these reports. The basic information needed for all reports includes: common name, age (adult, immature, etc.), plumage (breeding, non-breeding, etc.), number seen, location (county, nearest town, etc.), and date. The more unusual reports require additional information such as time seen, light conditions, optical equipment, distance to bird and how measured, as well as a description of the bird's size, shape, color pattern, vocalizations if any were heard, and behavior. It is also important to eliminate any similarly appearing species and discuss prior experience with both the species being reported and any with which it might be confused. The names and addresses of additional observers should be included as well as any reference material consulted and how it influenced the description. If it is possible to obtain photographs or audio recordings, be sure to include copies with the report. **Above all else, remember that the documentation process is in place to gather additional, substantiating information and is not meant to intimidate potential contributors or question the authenticity of their reports.**

Reports of birds should be submitted online via the society's website at [www.arbirds.org](http://www.arbirds.org) If further assistance is needed, contact the Curator at [arbirds@arbirds.org](mailto:arbirds@arbirds.org).

**Dedication**

This 2015 edition of the Arkansas Audubon Society field reference honors Dr. Douglas A. James. In the 1930s, when Dr. James was in grade school in suburban Detroit, he was often called upon to lead bird walks, using his 3X opera glasses and Chester Reed's Bird Guide (1931). He came to UA-Fayetteville in 1953, the start of a career that now includes guiding more than 130 MS and PhD graduate students. He is a founder of Arkansas Audubon Society (1955) and was its first curator of bird records. As President of the Wilson Ornithological Society (1977-79), he formally saluted the importance of amateurs in ornithology. His statewide collection of vetted bird observations, mostly submitted by amateurs, formed the core of Arkansas Birds (1986) and previous editions of this field list.

The Arkansas Audubon Society was formed in May 1955 to promote the protection and wise use of our natural resources. Weekend meetings are held twice each year at locations throughout the state and are generally scheduled to coincide with the peaks of spring and fall migration. Guest speakers, workshops, and guided field trips are featured at all meetings. A quarterly newsletter is published to keep the membership informed regarding current conservation issues, noteworthy bird records, and news of members and upcoming events. Membership is open to all interested persons. For further information, please visit the society's website at [www.arbirds.org](http://www.arbirds.org)

The 2015 Field Reference was compiled by Lyndal York. Document design and typesetting by Kenny Nichols and Kelly Chitwood. Additional editorial assistance provided by Leif Anderson, Dick Baxter, Rob Doster, Douglas A. James, Mike Mlodinow, Joseph C. Neal, LaDonna Nichols, Helen Parker, Herschel Raney, and Dan Scheiman.

To request 15 or more hardcopies of this field list, e-mail the Curator at [arbirds@arbirds.org](mailto:arbirds@arbirds.org) to make arrangements. A donation is appreciated to offset the costs of printing and mailing. To make a donation visit [www.arbirds.org](http://www.arbirds.org)

# Arkansas Audubon Society

## 2015 Field Reference to Arkansas Birds

Date \_\_\_\_\_ Time \_\_\_\_\_ Total Species Observed \_\_\_\_\_

Location \_\_\_\_\_ County \_\_\_\_\_

Observers \_\_\_\_\_

This publication provides information on 415 species of birds currently documented for Arkansas. It is based primarily on bird records maintained by the Arkansas Audubon Society, and secondarily on other sources, including *Arkansas Birds- Their Distribution and Abundance* (James and Neal, UA Press, 1986), Breeding Bird Surveys, Christmas Bird Counts, eBird, and published literature on Arkansas birds. **Since the information that follows is generalized due to space constraints, it may not seem appropriate for a given species in some locales.** Taxonomy and nomenclature follow the *Checklist of North American Birds*, Seventh Edition, AOU, 1998, as amended through the Fifty-sixth supplement, July 2015.

Approximate dates of arrival and departure are indicated by numbers 1-12 for the months of the year and letters A, B, C, D for the weeks of the month. A plus (+) sign immediately following indicates that discontinuous records outside the period are on file. For species that have nested in Arkansas, one of two symbols follows the name: an asterisk (\*) means the species currently nests with regularity, either yearly, or at least, with some consistency; an open circle (°) means that the species does not nest regularly and that there are few or no such reports since 1950. Beneath the dates of arrival and departure are codes which categorize seasonal status, relative abundance, and, in some cases, the sections of the state of primary occurrence.

The months of the year are grouped into the following seasons:  
**Sp**-spring (March-May)                      **S**-summer (June-July)  
**F**-fall (August-November)                      **W**-winter (December-February)

In combination with these designations, the following abbreviations are used to denote seasonal status:  
**P**-Present, species seen in the state each month but not necessarily in all sections (Great Egret)  
**M**-Migrant, species passes through the state in spring and/or fall (Cape May Warbler)  
**R**-Resident, species remains for an extended period of time (Painted Bunting)  
**L**-Local, species occurs in only a few places (Great-tailed Grackle)  
**V**-Visitor, species usually not present for an extended period of time (Black-legged Kittiwake)  
**Sa**-Storm-assisted, species appears in the state in association with strong coastal storms; not usually present for an extended time (Sooty Tern)

The categories of relative abundance assume reasonable effort spent in suitable habitat during appropriate seasons and are defined as follows:  
**c**-common, always or almost always observed (Killdeer)  
**fc**-fairly common, seen on a majority of trips but usually in smaller numbers (Red-shouldered Hawk)  
**u**-uncommon, may be difficult to find; seen only on a minority of trips (Golden-winged Warbler)  
**r**-rare, may be seen annually or nearly so but not in all locales (Golden Eagle)  
**vr**-very rare, encountered at irregular or infrequent intervals (Ruff)  
**occ**-occasional, 3-4 state records overall with at least 2 records since 1950 (Long-tailed Jaeger)  
**acc**-accidental, 1-2 state records (Fork-tailed Flycatcher)  
**irr**-irruptive, variable in occurrence, number, and distribution (Red-breasted Nuthatch)  
The abbreviations in parentheses following the statements of relative abundance indicate the primary but not necessarily the exclusive section of the state where the species occurs: **(n)** north, **(s)** south, **(e)** east, **(w)** west, **(ce)** central, **(nc)** north-central, **(sc)** south-central, **(ec)** east-central, **(wc)** west-central **(ne)** northeast, **(se)** southeast, **(nw)** northwest, and **(sw)** southwest.

Woodpecker, Pileated*	P fc	<b>Family Lanidae</b> <b>Shrikes</b>	
Ivory-billed	Formerly P Recent status unclear	Shrike, Loggerhead*	P SRu (Lfc e, nw) WRfc
<b>Order Falconiformes</b> <b>Falcons</b>		Northern	WVvr (5, 2011)
Caracara, Crested	SpVacc (1, 2015)	<b>Family Vireonidae</b> <b>Vireos</b>	
Kestrel, American*	P Mc SRu WRc	Vireo, White-eyed*	3C-11D+ SRc WVvr (s)
Merlin	9B-4C+ Mu WVu	Bell's*	4C-9D Mu SRu
Falcon, Peregrine°	8C-5B+ Mu WVr	Yellow-throated*	3D-10C+ SRfc
Prairie	10C-2B+ WVr	Blue-headed	9A-5D+ Mfc WRr (s)
<b>Order Psittaciformes</b> <b>Parakeets</b>		Warbling*	4B-9D+ SRLu Mfc
Parakeet, Carolina	extinct	Philadelphia	4C-5D 9B-10B+ Mfc
<b>Order Passeriformes</b> <b>Family Tyrannidae</b> <b>Flycatchers</b>		Red-eyed*	4A-10B+ SRc
Flycatcher, Olive-sided	4D-6D 8A-10B Mu	<b>Family Corvidae</b> <b>Jays and Crows</b>	
Wood-Pewee, Eastern*	4A-10D SRc	Jay, Blue*	P c
Flycatcher, Yellow-bellied	4D-5D 8B-10C SpMr FMvr	Nutcracker, Clark's	SpVacc (1, 1891)
Acadian*	4A-9D+ SRfc	Crow, American*	P c
Alder	5A-6A 8A-9B+ Mu	Fish*	P SRc WRLfc (s)
Willow*	5A-9C SpMu SRLr (nw)	<b>Family Alaudidae</b> <b>Larks</b>	
Least	4C-5D 8B-10B SpMfc FMu	Lark, Horned*	P WRc SRLu
Phoebe, Eastern*	P SRc SRr (se) WRfc	<b>Family Hirundinidae</b> <b>Swallows</b>	
Say's	9C-4D WVr	Martin, Purple*	2A-9B+ SRc
Flycatcher, Vermilion	10A-4B+ WVvr	Swallow, Tree*	2C-11C+ Mc SRLfc
Ash-throated	FWVacc (2, 2010)	Northern Rough-winged*	3A-10D+ SRc
Great-crested*	4B-10A+ SRc	Bank*	4A-5D 7A-9D+ Mfc SRvr
Kingbird, Couch's	WVacc (1, 1995)	Cliff*	3B-10B+ Mc SRc
Cassin's	SpSVacc (2, 1967)	Cave	3D-5D 6D-10A+ SFMr (sw, se) SpVvr
Western*	3D-10C Mu SRLu	Barn*	3A-11D+ SRc
Eastern*	3D-10D SRc	<b>Family Paridae</b> <b>Chickadees and Titmice</b>	
Gray	SpVacc (1, 2007)	Chickadee, Carolina*	P c
Flycatcher, Scissor-tailed*	3D-11D+ SRc	Titmouse, Tufted*	P c
Fork-tailed	WVacc (1, 1995)		

<b>Family Sittidae</b> <b>Nuthatches</b>		Thrush, Varied	WVacc (1, 2010)
Nuthatch, Red-breasted	9B-5B+ WRirr	<b>Family Mimidae</b> <b>Mockingbirds, Thrashers, and Allies</b>	
White-breasted*	P c (n, wc)	Catbird, Gray*	P SRc WRr
Brown-headed*	P c (s, wc)	Thrasher, Brown*	P c
<b>Family Certhiidae</b> <b>Creepers</b>		Sage	SpFVacc (2, 2004)
Creep, Brown°	10A-4C+ WRfc	Mockingbird, Northern*	P c
<b>Family Troglodytidae</b> <b>Wrens</b>		<b>Family Sturnidae</b> <b>Starlings</b>	
Wren, Rock	10B-4A+ WRr (w)	Starling, European*	P c
House*	P Mfc WRu SRfc (nw)	<b>Family Motacillidae</b> <b>Pipits</b>	
Winter	10A-4D+ WRfc	Pipit, American	9D-4C+ Mfc WRfc
Sedge*	P Mfc SRLfc (e, nw) WRLu	Sprague's	10A-4D Mr WRr
Marsh	9C-5C Mfc WRu	<b>Family Bombycillidae</b> <b>Waxwings</b>	
Carolina*	P c	Waxwing, Bohemian	1D-3C+ WVvr
Bewick's*	P SRr (nw) WRr	Cedar*	P SRu (nw) WRc
<b>Family Polioptilidae</b> <b>Gnatcatchers</b>		<b>Family Calcaridae</b> <b>Longspurs and Snow Bunting</b>	
Gnatcatcher, Blue-gray*	3C-10C+ SRc WVr	Longspur, Lapland	11B-3A+ WRfc
<b>Family Regulidae</b> <b>Kinglets</b>		Chestnut-collared	12D-1D+ WVvr
Kinglet, Golden-crowned	9D-4D WRc	Smith's	11A-3C Mr WRLu (e)
Ruby-crowned	9D-5A+ WRc	McCown's	FVacc (2, 1996)
<b>Family Turdidae</b> <b>Bluebirds, Thrushes, and Allies</b>		Bunting, Snow	11C-12D+ WVvr
Wheat, Northern	FVacc (1, 1990)	<b>Family Parulidae</b> <b>Wood Warblers</b>	
Bluebird, Eastern*	P c	Ovenbird*	4B-10B+ SRc (n, ce, wc)
Mountain	FSpVacc (2, 2013)	Warbler, Worm-eating*	4A-9D+ SRu
Solitaire, Townsend's	10C-4A WRvr (wc)	Waterthrush, Louisiana*	3C-10B SRfc
Veery	4C-5C+ SpMu	Northern	4B-5C 8B-10A+ SpMc FMfc
Thrush, Gray-cheeked	4A-5D 9B-10C+ SpMfc FMvr	Warbler, Bachman's	extinct
Swainson's	4A-5D 9A-10C+ SpMc FMu	Golden-winged	4D-5B 8D-10B Mu
Hermit	10A-4D+ WRc	Blue-winged*	4A-10B Mfc SRLu (n, ce)
Wood*	4A-10B+ SRfc	Black-and-white*	3C-10B+ SRc WVvr (s)
Robin, American*	P c	Prothonotary*	4A-9C+ SRfc

Warbler, Swainson's*	4A-9B+ SRLu	<b>Family Emberizidae</b> <b>New World Sparrows</b>	
Tennessee	4B-5C 9A-10C+ Mc	Towhee, Green-tailed	FVacc (1, 1952)
Orange-crowned	9D-5C+ Mfc WRu	Spotted	10C-4A+ WRr
Nashville	4A-5C 9A-10D+ Mc	Eastern*	P SRfc WRc
Connecticut	4C-5D SpMvr	Sparrow, Rufous-crowned*	P Lr (we)
Mourning	4D-5D 8D-9D+ Mu	Cassin's°	SpFVacc (2, 2011)
Kentucky*	4B-9D+ SRfc	Bachman's*	4A-9C+ SRLu
Yellowthroat, Common*	4A-10D+ SRc WRu (s, e)	American Tree	10B-4A Mr WRr (n)
Warbler, Hooded*	4A-10A+ SRfc	Chipping*	P SRc (n, wc) WRc (s)
Redstart, American*	4B-10C Mfc SRu (Lfc nw)	Clay-colored	4B-5B 9A-11B+ Mu (w) Mr (e)
Warbler, Cape May	4D-5C 8D-11D+ Mr	Brewer's	WVacc (1, 2011)
Cerulean*	4C-9A+ SRu (Lfc nw)	Field*	P SRfc WRc
Parula, Northern*	3C-10C+ SRc	Vesper	10A-4D Mfc WRu (s, e)
Warbler, Magnolia	4B-5D+ 9A-10B+ Mfc Mc (e)	Lark*	3D-9D+ SRfc SRu (nw) WVvr (s)
Bay-breasted	4B-6B 9D-10B+ Mu	Bunting, Lark	WSpVacc (3, 2013)
Blackburnian	4D-6A 9A-10B Mu Mfc (e)	Sparrow, Savannah	10A-5C+ WRc
Yellow*	4B-9D+ Mfc SRLu (nw)	Grasshopper*	4A-11D+ SRLu SRLfc (nw)
Chestnut-sided*	4D-10B+ SpMfc FMfc (e) u (w) SRLr (nc, nw)	Henslow's*	P Mr SRLvr (nw, n, wc) WRr (se)
Blackpoll	4B-5D SpMfc	Le Conte's	10A-5B WRLfc
Black-throated Blue	4C-5B 9C-10C+ Mvr	Nelson's	5B-5C 9D-10D+ Mr
Palm	4C-5A 9C-11C+ Mfc WRr	Fox	10B-4C+ WRfc
Pine*	P c WRu (nw)	Song	10A-5B+ SRc
Yellow-rumped	9D-5B+ WRc	Lincoln's	10A-5C+ Mfc (w, ce) WRu (w)
Yellow-throated*	3D-10C+ SRfc	Swamp	10A-5A+ WRc
Prairie*	4B-9C+ SRfc	White-throated	10A-5B+ WRc
Black-throated Gray	FVacc (1, 2005)	Harris's	10C-5B WRu (w, ce)
Black-throated Green*	3D-10C+ Mc SRLu (nc, nw)	White-crowned	10A-5D WRc
Canada	4D-6A 8D-10A Mu	Junco, Dark-eyed	10B-4B+ WRc
Wilson's	4D-6A 8D-10A+ Mu (e) Mc (w)	<b>Family Cardinalidae</b> <b>Tanagers, Cardinal, and Grosbeaks</b>	
Chat, Yellow-breasted*	4C-9D+ SRc	Tanager, Summer*	4B-10B+ SRc WVvr

Tanager, Scarlet*	4B-10B+ Mfc SRfc (n, wc)	Crossbill, Red°	9A-7D+ Vr
Western	8D-5B WVvr	White-winged	1A-4C WSpVvr
Cardinal, Northern*	P c	Redpoll, Common	12A-4A WVvr (n)
Grosbeak, Rose-breasted	4C-5C 9C-10C+ SpMfc FMu	Siskin, Pine°	10B-5B+ Mrr WRirr
Black-headed	9D-5B+ WVvr	Goldfinch, Lesser	SpSWVacc (4, 2012)
Blue*	4B-10B SRc	American*	P WRc SRc (nw, ce)
Bunting, Lazuli	2D-6B SpVvr	Grosbeak, Evening	11B-5A+ Wrr vr
Indigo*	4B-11A+ SRc WVvr	<b>Family Passeridae</b> <b>Old World Sparrows</b>	
Painted*	4B-9D+ SRfc	Sparrow, House*	P c
Dickcissel*	4D-10B+ SRc WVvr		
<b>Family Icteridae</b> <b>Blackbirds and Allies</b>			
Bobolink	4C-5D 8C-10B+ SpMfc FMLu		
Blackbird, Red-winged*	P c		
Meadowlark, Eastern*	P c		
Western	10C-4D+ WRu		
Blackbird, Yellow-headed	4A-5C+ 8C-10C+ SpMu FMr WVvr		
Rusty	10C-4C+ WRfc		
Brewer's	11A-4B+ WRfc (e, sw) WRu (nw)		
Grackle, Common*	P c		
Great-tailed*	P Lu (w)		
Cowbird, Brown-headed*	P c		
Oriole, Orchard*	4A-9C+ SRc		
Bullock's	12D-2C WVvr		
Baltimore*	4A-9D+ SRLfc (ce, e, s) SRLu (nw) WVvr (s)		
Scott's	WSpVacc (3, 2013)		
<b>Family Fringillidae</b> <b>Finches</b>			
Rosy-Finch, Gray-crowned	SpVacc (1, 2012)		
Grosbeak, Pine	WSpSVacc (4, 1989)		
Finch, House*	P c		
Purple°	10D-4B+ WRfc		